

Asteroseismic constraints on the OPAL opacity interpolation

W. M. Yang and M. Li

Department of Physics and Chemistry, Henan Polytechnic University, Jiaozuo 454000, China
 email: yang.wuming@yahoo.com.cn

Abstract. The frequency difference between a model used only two-point interpolation of opacity and a model used piecewise linear interpolation of opacity is of the order of several microHertz at a certain stage, which is almost 10 times worse than the observational precision of p-modes of solar-like stars. Therefore, the two-point interpolation of opacity is unsuitable in modelling of solar-like stars with element diffusion.

Keywords. stars: evolution – diffusion – stars: oscillations

1. Introduction

Timescales of evolution and element diffusion are similar in solar-like stars (Turcotte et al. 1998). Thus element diffusion should be calculated in modelling of solar-like stars. Furthermore, helioseismology has confirmed the importance of including element diffusion and settling in solar modelling (Christensen-Dalsgaard et al. 1993; Basu et al. 2000).

Heavy-element abundance, Z , is constant in main-sequence (MS) stage of solar-like stars without considering metal settling, thus evolution of these stars need only one set of opacity tables at a fixed Z_1 . However the Z is a variable in the models with metal settling and then the second set of opacity tables must be obtained at a fixed Z_2 . Opacity at the desired Z , X , T , and ρ can be obtained by two-point interpolation, i.e.,

$$\kappa(Z, X, T, \rho) = \kappa_2(Z_2, X, T, \rho) + \frac{\kappa_1(Z_1, X, T, \rho) - \kappa_2(Z_2, X, T, \rho)}{Z_1 - Z_2} \cdot (Z - Z_2). \quad (1.1)$$

2. Calculation and results

In order to study the impact of opacity interpolation, using the Yale Rotating Evolution Code (YREC7) in its nonrotating configuration, we construct four models listed in Table 1. All parameters of the models are same except the parameters of opacity interpolation. All models evolve from pre-main sequence (PMS) to somewhere near the end of the MS. The OPAL eos (Rogers & Nayfonov 2002), OPAL opacity (Iglesias & Rogers 1996), and the Alexander & Ferguson(1994) opacity for low temperature are used. Element diffusion is implemented following the prescription of Thoul et al.(1994).

The changes in the effective temperature between MT2, MT3, and MM1 at the same age are a few Kelvin, which is within the error of observation of stellar effective temperature. The frequency differences between MT1 and MT2 are zero. In Fig. 1A, we represent the frequency differences between MT2 and MT3 at the same age. The difference increases from about $1 \mu\text{Hz}$ at the age of 1 Gyr to around $4 \mu\text{Hz}$ at the age of 6 Gyr. But the difference at the age of 7 Gyr is less than that at the age of 6 Gyr. The frequency differences between MM1 and MT3 are shown in Fig. 1B. The differences arrive at a maximum at the age of about 3 Gyr. Then with increase in age, the differences

Table 1. Model parameters.

Model	Mass $M_{\odot}$	Z_0	Two-point interpolation Z_1 — Z_2	Piecewise linear interpolation $\delta z = Z_{i+1} - Z_i$
MT1	1.10	0.022	0.022 — 0.021	
MT2	1.10	0.022	0.022 — 0.020	
MT3	1.10	0.022	0.022 — 0.019	
MM1	1.10	0.022		0.001

Notes: The Z_0 is the initial metal abundance. Piecewise linear interpolation: if $Z_i \leq Z < Z_{i+1}$, YREC7 will interpolate between $\kappa(Z_i)$ and $\kappa(Z_{i+1})$ to obtain the opacity at the required Z .




Figure 1. Frequency differences at different ages labeled by the different symbols. A: $\nu_{MT2}-\nu_{MT3}$; B: $\nu_{MM1}-\nu_{MT3}$; C and D: $\nu_{MM1}-\nu_{MT2}$.

decrease. At the age of around 6 Gyr, the differences are almost zero. Thus the discrepancy between MM1 and MT3 mainly occurs between the ages of 2 and 4 Gyr. In Fig. 1C and D, we represent the frequency differences between MM1 and MT2. The differences are almost zero when the age of the models is less than 3 Gyr. Then with increase in age, the frequency differences increase. At the age of around 6 Gyr, the differences arrive at a maximum. The differences at the age of 4 - 7 Gyr are of the order of several microHertz, which is almost 10 times larger than the uncertainty of observation of stellar p-modes that is expected to reach 0.1-0.4 μHz (Thèdo et al. 2005; Bedding et al. 2004).

3. Discussion

Heavy-element abundance is a constant in the models of PMS, and the Z_s and Z_c are close to the Z_0 in the early evolutionary stage of MS. Thus Z_1 is specified to be Z_0 in models MT1, MT2, and MT3, and one of Z_i should be equal to Z_0 in MM1. The differences of the opacity interpolation in MT2, MT3, and MM1 should result in the difference between MT2, MT3, and MM1 and then lead the frequency difference between MT2, MT3, and MM1. The frequency difference between the models MT2, MT3, and MM1 is of the order of several microHertz at a certain phase, which is almost 10 times worse than the observational precision of p-modes of solar-like stars. Consequently, in modelling of solar-like star with metal settling, the two-point interpolation of opacity is unsuitable, and at least piecewise linear interpolation is required.

References

- Alexander, D. R., & Ferguson, J. W. 1994, *ApJ*, 437, 846
 Basu, S., Pinsonneault, M. H., & Bahcall, J. N. 2000, *ApJ*, 529, 1084
 Bedding, T. R., Kjeldsen, H., Butler, P. R., et al. 2004, *ApJ*, 614, 380
 Christensen-Dalsgaard, J., Proffitt, C. R., & Thompson, M. J. 1993, *ApJ*, 403, L75
 Iglesias, C., & Rogers, F. J., 1996, *ApJ*, 464, 943
 Rogers, F. J., & Nayfonov, A., 2002, *ApJ*, 576, 1064
 Thado, S., Vauclair, S., Castro, M., Charpinet, S., & Dolez, N. 2005, *A&A*, 437, 553
 Thoul, A. A., Bahcall, J. N., & Loeb, A. 1994, *ApJ*, 421, 828
 Turcotte, S., Richer, J., & Michard, G. 1998, *ApJ*, 504, 559